

## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

DAA/Goddard

(NASA-CR-176142) LOW-EXCITATION HERBIG-HARO  
OBJECTS AND INTERSTELLAR EXTINCTION Final  
Technical Report, 1 Aug. 1984 - 31 Jul. 1985  
(Missouri Univ.) 2 p HC A02/MF A01 CSCL 03B

N85-34726

Unclass  
63/90 22131

Low-Excitation Herbig-Haro Objects and Interstellar Extinction

Final Technical Report

Richard D. Schwartz

August 1, 1984-July 31, 1985

Physics Department

University of Missouri, St. Louis

801 Natural Bridge Road

St. Louis, MO 63121

NASA NAG 5-461



Previous work on low-excitation Herbig-Haro (HH) objects (Schwartz 1983)<sup>1</sup> has confirmed the existence of a strongly-enhanced hydrogen two-photon continuum which is produced by low-velocity shocks in neutral material. In the present study, we have compared the theoretical two-photon energy distribution with that observed for HH 43. With the addition of optical ( $3300 < \lambda < 7000\text{\AA}$ ) data to the ultraviolet ( $1200 < \lambda < 3100\text{\AA}$ ) data obtained with the I.U.E, the wavelength dependence of interstellar extinction toward HH 43 has been determined.

In summary, the extinction curve appears to follow that found for the nearby  $\theta$  Ori group of stars, with nearly neutral UV extinction, and an enhanced ratio of total to selective absorption ( $A_V/E(B-V) > 5$ ). A best fit of observed data to the theoretical curve yields  $E(B-V) = 0.2$ . Continuum and emission-line data for HH 43 have been corrected for extinction, and an analysis of the energetics of the shock-excited pockets of gas has been carried out. Evidence suggests that a young, embedded stellar object with a luminosity of about  $5 L_\odot$  is producing a collimated, supersonic flow of material in which the HH 43 shocks appear with a luminosity of  $\sim 0.25 L_\odot$ . The mechanical luminosity associated with the outflow must therefore be a substantial fraction of the total luminosity of the young star.

This work has been reported at the January, 1985 meeting of the A.A.S. in Tucson (Schwartz and Dopita 1984), and a detailed paper is in press (Schwartz, Dopita, and Cohen 1985).

---

<sup>1</sup>Work carried out with support of NASA grant NAG 5-243.

#### References

- Schwartz, R. D. 1983, Ap. J. Lett. 268, L37.  
Schwartz, R. D. and Dopita, M. A. 1984. B.A.A.S. 16, 972.  
Schwartz, R. D., Dopita, M. A., and Cohen, M. 1985.  
Astron. J. (September, in press).